PED Low Titer O+ Whole Blood

History

- What was the mechanism of injury blunt (MVC, fall, blow to body) vs. penetrating (stabbing, GSW, foreign body)?
- Did a medical condition contribute to the mechanism of injury? Other medical conditions?
- Medications Coumadin? Plavix? Aspirin? Pradaxa? Xarelto? Eliquis? (any blood thinners or anticoagulants)

Key Concepts

 Low Titer O + Whole Blood is now being used to treat severely injured trauma patients who have or are at risk for severe hemorrhage

MARCHES Protocol

- Massive bleeding control
- Airway NPA/OPA/ Crich
- Respiratory decompress chest if tension pneumothorax, occlusive dressing for open pneumothoraces
- Circulation- IV/IO Tgx, pelvic binder, wound packing
- Hypothermia care
- Eye injuries cover with rigid shield and no pressure on the eye
- Spinal motion restriction if indicated

Medical Patients in Hemorrhagic Shock

 Less likely for children but consider the same medical causes for hemorrhagic shock as in adults

Criteria

HEMORRHAGIC SHOCK in medical or trauma Adult and Pediatric (≥ 6 yo) patients

Relative Contraindications

- Patient < 6 years old
 - Consult Medical Direction if patient is in hemorrhagic shock and < 6 yo
 - Medical Director may elect to give blood in this patient population
 - o Religious objection to receiving whole blood—consult On Call Medical Director

EMT

- Follow Trauma General Patient Care Guideline
- Follow appropriate Trauma Guideline

Paramedic

For Patients in HEMORRHAGIC SHOCK:

Administer Whole Blood with signs of acute hemorrhagic shock as evidenced by:

- Systolic blood pressure < 70 mmHg OR
- Systolic Blood Pressure < 90 mmHg with Heart Rate ≥ 110 beats per min OR
- ETCO2 < 25 OR
- Witnessed arrest from hemorrhagic shock < 5 min prior to provider arrival and continuous CPR throughout downtime

In general only 500mL (1 unit) of Low Titer O+ Whole Blood (LTO+WB) will be available per patient. If more than 500 mL of Whole Blood is available on scene the following general guidelines apply:

- 5-10 yo are eligible for 500 mL of Whole Blood
 - o Consult Medical Direction for further orders, if needed
- 11-13 yo are eligible for 1000 mL of Whole Blood
 - o Consult Medical Direction for further orders, if needed
- ≥13 yo are eligible for >1000 mL of Whole Blood
 - Consult Medical Direction for further orders, if needed

Of Note: At this time the LTO+WB does not have unit markings on the bag.